

Colloid bonded Medical Compounds

Edith Malecki, Helmut Rosemeyer

Europäische Patentanmeldung EP 12186555.4; 28.09.2012

Colloid bonded Medicinal Compounds

The invention relates to colloids bound medicinal compounds or fluorescent markers, to a process for the preparation thereof, and to a pharmaceutical formulation containing such compounds.

Achieving good solubility and/or chemical and metabolic stability can be a challenging issue in drug development. As a result a significant number of drugs or drug candidates exhibit sub-optimal properties potentially resulting in repeated administration, high doses and associated cost of goods, and represent a burden both to the patient and the drug companies developing them. Over the last 30 years a few innovative approaches have been developed to try to address these issues either via formulation strategies and/or bioconjugation methodologies such as PEGylation. Starch derivatives (eg HES) are water soluble semi-synthetic polysaccharides that have been used in large amounts for decades in hospital care notably for the treatment of hypovolemia and that exhibit low toxicology potential. Their behavior in the human body is therefore very well understood. It has recently been reported that they could also be used as a biopolymer conjugates and could represent as such an elegant alternative to other methodologies described above.

The covalent binding to colloids enables substances to be introduced by phagocytosis into cells of the immune system, which would not be taken up, or if so only in negligible amounts, without such modification. EP 1 230 935 A1 describes the chemical binding of medicinally active substances to a polysaccharide to form a linker. The uptake of substances by correspondingly specialized cells of the reticulohistiocytic system has been demonstrated for a wide variety of colloids and particles. However, there is still a need to improve the binding complexes in terms of a sufficient availability of the medicinally active substance and/or the fluorescence marker at the cells which also requires a effective enzymatic cleavage of the

medicinally active substance and/or the fluorescence marker from the colloid active substance.


For the introduction of medicinally active substances or fluorescence marker into specific organs and cell systems of the body, the following conditions must be met:

1. The medicament complex, which consists of the medicament chemically bonded to a colloid, should be water-soluble and circulate in the blood for a sufficient period of time.
2. The medicament complex should have no influence on blood clotting.
3. The medicinally active substance must be cleavable from the colloid active substance, especially enzymatically cleavable.

The object of the present invention is to provide a novel and versatile way to link colloids, especially HES molecules or derivatives, to an active ingredient or a fluorescence marker via a phosphor acid diester linkage therefore giving access to novel bioconjugates. These bioconjugates retain biological activity while exhibiting improved drug properties.

Surprisingly, it has now been found that bonding of a medicinally active substance or a fluorescence marker to a colloid (colloid-active compound) via a phosphor acid diester solves the above mentioned problems and serves, in particular, as a suitable transport system for medicaments and/or fluorescence markers covalently linked thereto.

The present invention relates to a compound of general formula (I)



wherein

X is a colloid-active compound;

L^1 is a first linker by means of which X and the phosphate group are covalently linked together;

L^2 is a second linker by means of which the phosphate group and A are covalently linked together;

A is a medically active substance or a fluorescence marker;

n is an integer of at least 1.

In a preferred embodiment of the present invention, n is an integer of from 1 to 10,000, preferably from 2 to 1000, more preferably from 5 to 500, especially from 10 to 100.

In a further preferred embodiment the colloid-active compound X is selected from the group consisting of amyloses, amylopectins, acemannans, arabinogalactans, galactomannans, galactoglucomannans, xanthans, carrageenan, hyaluronic acid, deacetylated hyaluronic acid, chitosan, starch and modified starch.

The modified starch is preferably selected from the group consisting of hydroxyalkyl starches, esterified starches, carboxyalkyl starches, hydroxyalkyl carboxyalkyl starch, aminated hydroxyalkyl starch, aminated hydroxyalkyl carboxyalkyl starch and aminated carboxyalkyl starch.

According to a especially preferred embodiment the modified starch is selected from hydroxyethyl starch or aminated hydroxyethyl starch. Preferably the degree of substitution, DS, of the modified starch, especially hydroxyethyl starch, is from 0.2 to 0.8, preferably from 0.3 to 0.6.


Advantageously, the colloid-active compound has an average molecular weight of from 20,000 to 800,000 daltons, preferably from 25,000 to 500,000 daltons, especially from 30,000 to 200,000 daltons.



According to a further preferred embodiment of the invention the medically

active compound A is selected from the group consisting of antibiotics, chemotherapeutics, cytostatic agents, antigens, oligonucleotides, mediators, false metabolic substrates, analgetics, antifungals and cytotoxic substances.

A in formula (I) is preferably selected from group consisting of 5-Fluorouracil, Amphotericin B, Daunorubicin, Doxorubicin, Cladribine, Zosyn (Tazosyn), Floxuridine, Delafloxacin, Gemcitabine, Irinotecan, Vadimezan, Fluconazole, Voriconazole and Ravuconazole.

Specifically said medicinally active compound A is selected from formula (II) to (IV)





wherein

R^1 is H or an organic moiety, preferably a C_1 - C_{28} chain which may be branched or linear and which may be saturated or unsaturated and which may optionally be interrupted and/or substituted by one or more hetero atom(s) (Het1) and/or functional group(s)(G1); or

R^1 is a C_3 - C_{28} moiety which comprises at least one cyclic structure and which may be saturated or unsaturated and which may optionally be interrupted

and/or substituted by one or more hetero atom(s) (Het1) and functional group(s)(G1);

R² is H or an organic moiety comprising 1 to 30 carbon atoms;

R³ and R⁴ represent independently from each other H or a C₁-C₂₈-alkyl moiety which may optionally be substituted or interrupted by one or more heteroatom(s) and/or functional group(s); or

R³ and R⁴ form a ring having at least 5 members, preferably a ring having 5 to 8 carbon atoms and wherein the ring may be substituted or interrupted by one or more hetero atom(s) and/or functional group(s);

R⁵ and R⁶ represent independently from each other H or a C₁-C₂₈-alkyl moiety which may optionally be substituted or interrupted by one or more heteroatom(s) and/or functional group(s);

or


R⁵ and R⁶ form a ring having at least 5 members, preferably a ring having 5 to 18 carbon atoms and wherein the ring may be substituted or interrupted by one or more hetero atom(s) and/or functional group(s);

R⁷ is a hydrogen atom or -O-R⁸;

R⁸ is H or C₁-C₂₈ chain which may be branched or linear and which may be saturated or unsaturated and which may optionally be interrupted and/or substituted by one or more hetero atom(s) (Het1) and/or functional group(s)(G1).


In a preferred embodiment substituent R¹ is a linear or branched chain comprising 1 to 50 carbon which may be interrupted and/or substituted by one or more hetero atom(s) (Het1) and/or functional group(s)(G1). Preferably, R¹ is a linear or branched chain comprising 2 to 40, more preferably 3 to 30, especially 4 to 28 or 6 to 20 or 8 to 16 carbon atoms. In one aspect of the invention R¹ is a linear or branched C₁-C₂₈-alkyl, preferably C₂-C₂₀-alkyl, more preferably C₄-C₂₀-alkyl or C₆-C₁₈-alkyl, especially C₈-C₁₆-alkyl which may be substituted or unsubstituted. In a further aspect of the invention the carbon chain is interrupted by one or more hetero atom(s) (Het1) wherein the Het1 is preferably selected from O, S and N, more preferably selected from O or N. In one aspect the substituent

R¹ is interrupted by up to 3 hetero atom(s) (Het1), preferably 1 or 2 hetero atoms such as O. In a further aspect of the invention the carbon chain of substituent R¹ is interrupted by nitrogen which preferably further branches the chain. An exemplary embodiment of this type of substituent is reflected in the following formula:




wherein R⁹ and R¹⁰ are independently selected from a C₁ to C₃₀ chain which can be saturated or unsaturated, preferably a C₁ to C₃₀ alkyl, preferably C₄ to C₂₄ alkyl, more preferably C₈ to C₂₂ alkyl and especially C₁₂ to C₁₈ alkyl; or a C₂ to C₃₀ chain having one or more carbon-carbon double and/or carbon-carbon triple bond(s); and

"a" is an integer ranging from 1 to 20, preferably 2 to 18, more preferably 3 to 12 or 4 to 8. However, the linking moiety which links the nitrogen atom with substituents R⁹ and R¹⁰ to the 5-fluorouracil moiety can also be a unsaturated carbon chain having one 2 to 20 carbon atoms and one or more carbon- carbon double and or carbon-carbon triple bonds. The exemplary substituent of the following formula:



can be synthesized by various synthetic routes. Scheme 1 shows several synthetic routes for precursors which can be attached to the 5-fluorouracil moiety.



Scheme 1

As can be seen from Scheme 1 various precursor for the connection with the nitrogen atom of the 5-fluorouracil moiety can be obtained by different synthetic routes. In a preferred embodiment of the present invention substituent R^1 is a double chained substituent. The double chained substituents can be obtained as reflected in Scheme 2.

In a first aspect the dioctadecylamine is reacted with methyl bromoacetate in the presence of dibenzo-[18]-crown-6 which leads to the pure methyl ester in almost quantitative yield. The ester can be reduced with LiAlH_4 to give the alcohol.


In order to extend the spacer between the hydroxyl group and the nitrogen carrying the carbon chains the dioctadecylamine can be reacted with methyl

acrylate which results in almost quantitative yield to the ester which was further reduced with LiAlH_4 to give a lipophilic aminopropanol derivative.

In a further aspect the dioctadecylamine was reacted with succinic anhydride to give the acid which can be converted to the methyl ester by reaction with dimethyl sulphate in the presence of K_2CO_3 . The methyl ester can then be reduced with LiAlH_4 yielding the further extended alcohol, namely a lipophilized 4-aminobutanol derivative.

In a further reaction the dioctadecylamine can be alkylated with 1,4-dichlorobut-2-yne in the presence of Na_2CO_3 in benzene.

In the following Scheme 2 various synthetic routes to obtain single chain precursor or double chain precursor with different chains for the substitution of the 5-fluorouracil moiety are disclosed. The single chain precursor reflected in Scheme 3 is interrupted by a hetero atom (N) or a functional group (amid; NHCO).



Scheme 2

As can be seen from Scheme 2 lipid single chain precursors can be obtained by the reaction of octadecylamine with succinic anhydride which leads to the acid which can be reduced with LiAlH_4 in THF at ambient temperature which leads to the reduction of the carboxylic group only, but not of the amide moiety and results into the amidoalcohol in 82% yield. Replacement of THF by Et_2O however results in the amino alcohol in a high yield of 84%. Subsequent reaction of amino alcohol with propargyl bromide results in the double chained alkyne in 61% yield.

It has surprisingly been found that the lipophilic carbon chains comprising a hydroxyl functional group or a halide can be introduced regioselectively into the 5-fluorouracil derivative. The lipophilic groups can principally be positioned either at the heterocyclic base or at the glyconic moiety and can be introduced

by various methods, e.g. by base-catalysed alkylation with alkyl halides.

The reaction of unprotected 5-fluorouracil derivatives with halogenated alkyls, alkenes or alkynes can be performed in DMF/K₂CO₃ (direct alkylation) and leads to the alkylation of the unsubstituted nitrogen atom in the 5-fluorouracil ring.

Preferably, the unsubstituted nitrogen atom in the 5-fluorouracil ring is substituted by an halogen substituted precursor under the proviso that the hydroxyl groups present in the 5-fluorouracil derivative are protected by protecting groups. Suitable protecting groups are known to the person skilled in the art. Examples are dimethoxytrityl (DMT) and a *tert*-butyl-dimethylsilyl group.

Surprisingly it has been found that the hydroxyl functional lipophilic precursor (such as the amino alcohols reflected in Scheme 1 and 2) can be selectively reacted with the unsubstituted nitrogen atom of the 5-fluorouracil derivative by a Mitsunobu reaction. This reaction is carried out by first protecting any hydroxyl groups which may be present in the 5-fluorouracil derivative.

The Mitsunobu reaction is generally carried out by reacting the alcohol and the 5-fluorouracil derivative which comprises the unsubstituted ring nitrogen atom in the presence of triphenylphosphine and diisopropylazodicarboxylate (DIAD).

Further, R¹ is preferably a C₂ to C₄₀ chain which is unsaturated, more preferably a C₈ to C₂₈ chain which is unsaturated. In one embodiment of the invention R¹ comprises one or more carbon-carbon double bond(s) and/or one or more carbon-carbon triple bond(s). In a particular preferred embodiment R¹ comprises two or more, especially 2 to 6, such as 2 to 4 carbon-carbon double bonds.

In a specially preferred embodiment the substituents are derived from nature. Suitable naturally derived substituents have a structure derived from terpenes. When terpenes are chemically modified such as by oxidation or rearrangement of the carbon skeleton, the resulting compounds are generally referred to as

terpenoids. In a preferred embodiment R^1 is a cyclic or alicyclic terpenoid, preferably a terpenoid having 8 to 36 carbon atoms.

The terpenes are preferably selected from monoterpenes, sesquiterpenes, diterpenes, sesterterpenes, triterpenes and sesquaterpenes.

Suitable monoterpenes or monoterpenoids which can be acyclic or cyclic are selected from the group consisting of geraniol, limonene, pinen, bornylen, nerol.

Suitable sesquiterpenes sesquiterpenoids which can be acyclic or cyclic may inter alia be selected from farnesol.

Suitable sesterterpenes or sesterterpenoids are inter alia selected from geranyl-farnesol.


Suitable diterpenes or diterpenoids can be selected from the group consisting of abietic acid, aphidicolin, cafestol, cembrene, ferruginol, forskolin, guanacastepene A, kahweol, labdane, lagochilin, sclarene, stemarene, steviol, taxadiene (precursor of taxol), tiamulin, geranylgeraniol and phytol.

According to a especially preferred embodiment of the invention R^1 is selected from the group consisting of geranyl, farnesyl, neryl and phythyl.

According to a further alternative aspect R^1 is H or C_3 - C_{28} chain which may be branched or linear and which may be saturated or unsaturated and which may optionally be interrupted and/or substituted by one or more hetero atom(s) (Het1) and/or functional group(s)(G1); or

R^1 is a C_1 - C_{28} moiety which comprises at least one cyclic structure and which may be saturated or unsaturated and which may optionally be interrupted and/or substituted by one or more hetero atom(s) (Het1) and functional group(s)(G1);

According to an especially preferred embodiment R^1 in formula (II) is selected from H,



substituted or unsubstituted cyclic terpene moieties,
wherein

R^9 and R^{10} are independently selected from C_1 to C_{30} alkyl,
 b is an integer ranging 1 to 4, preferably n is 1 or 2; and
 a is an integer ranging from 1 to 20, preferably 2 to 18.

In a further aspect of the invention group A is a fluorescence marker which is preferably selected from the group consisting of fluorescein isothiocyanate (FITC), phycoerythrin, rhodamide, 2-aminopyridine and coumarine dyes.

In a further aspect of the invention linker L^1 and/or L^2 are independently selected from a single bond or a saturated or unsaturated moiety having 1 to 30, preferably 2 to 20 carbon atoms, more preferably a carbon chain which may be substituted and/or interrupted by one or more functional groups selected from carboxylic acid ester, phosphate ester, carboxylic acid amides, urethane, ether and amine groups. L^1 and/or L^2 may also comprise cyclic moieties.

According to a preferred embodiment linker L^1 and/or L^2 are independently selected from a single bond; alkandiyyl, preferably C_1 - C_{20} -alkandiyyl; alkendiyyl, preferably a C_2 - C_{20} -alkendiyyl; alkyndiyyl, preferably a C_2 - C_{20} -alkyndiyyl; aryl moiety, aralkyl moiety and heterocyclic moiety.

Preferably, the alkandiyl represents a straight-chain or branched-chain alkandiyl group bound by two different carbon atoms to the molecule, it preferably represents a straight-chain or branched-chain C₁₋₁₂ alkandiyl, particularly preferably represents a straight-chain or branched-chain C₁₋₆ alkandiyl; for example, methandiyl (--CH₂--), 1,2-ethandiyl (--CH₂-CH₂--), 1,1-ethandiyl ((--CH(CH₃)--), 1,1-, 1,2-, 1,3-propanediyl and 1,1-, 1,2-, 1,3-, 1,4-butanediyl, with particular preference given to methandiyl, 1,1-ethandiyl, 1,2-ethandiyl, 1,3-propanediyl, 1,4-butanediyl.

Further, preferably the alkendiyl represents a straight-chain or branched-chain alkendiyl group bound by two different carbon atoms to the molecule, it preferably represents a straight-chain or branched-chain C₂₋₆ alkendiyl; for example, --CH=CH--, --CH=C(CH₃)--, --CH=CH-CH₂--, --C(CH₃)=CH-CH₂--, --CH=C(CH₃)--CH₂--, --CH=CH-C(CH₃)H--, --CH=CH-CH=CH--, --C(CH₃)=CH-CH=CH--, --CH=C(CH₃)-CH=CH--, with particular preference given to --CH=CH-CH₂--, --CH=CH-CH=CH--.

The aryl moiety preferably represents an aromatic hydrocarbon group, preferably a C₆₋₁₀ aromatic hydrocarbon group; for example phenyl, naphthyl, especially phenyl which may optionally be substituted. The aromatic moiety may form a linker in that the aromatic moiety is attached to the phosphor acid diester and the group A. The phosphor acid ester group and group A may be in ortho, meta or preferably para-position of the aromatic moiety.

Aralkyl moiety denotes an "Aryl" bound to an "Alkyl" and represents, for example benzyl, α-methylbenzyl, 2-phenylethyl, α,α-dimethylbenzyl, especially benzyl. Aralkyl moiety can be attached to the phosphoric acid ester group via the alkyl or via the aryl part of the aralkyl moiety. Likewise group A can be attached to the phosphoric acid ester group via the alkyl or via the aryl part of the aralkyl moiety.

Heterocyclic moiety represents a saturated, partly saturated or aromatic ring

system containing at least one hetero atom. Preferably, heterocycles consist of 3 to 11 ring atoms of which 1-3 ring atoms are hetero atoms. Heterocycles may be present as a single ring system or as bicyclic or tricyclic ring systems; preferably as single ring system or as benz-annelated ring system. Bicyclic or tricyclic ring systems may be formed by annelation of two or more rings, by a bridging atom, e.g. oxygen, sulfur, nitrogen or by a bridging group, e.g. alkandiyl or alkenediyl. A Heterocycle may be substituted by one or more substituents selected from the group consisting of oxo ($=O$), halogen, nitro, cyano, alkyl, alkoxy, alkoxyalkyl, alkoxycarbonyl, alkoxycarbonylalkyl, halogenalkyl, aryl, aryloxy, arylalkyl. Examples of heterocyclic moieties are: pyrrole, pyrroline, pyrrolidine, pyrazole, pyrazoline, pyrazolidine, imidazole, imidazoline, imidazolidine, triazole, triazoline, triazolidine, tetrazole, furane, dihydrofurane, tetrahydrofurane, furazane (oxadiazole), dioxolane, thiophene, dihydrothiophene, tetrahydrothiophene, oxazole, oxazoline, oxazolidine, isoxazole, isoxazoline, isoxazolidine, thiazole, thiazoline, thiazolidine, isothiazole, isothiazoline, isothiazolidine, thiadiazole, thiadiazoline, thiadiazolidine, pyridine, piperidine, pyridazine, pyrazine, piperazine, triazine, pyrane, tetrahydropyrane, thiopyrane, tetrahydrothiopyrane, oxazine, thiazine, dioxine, morpholine, purine, pterine, and the corresponding benz-annelated heterocycles, e.g. indole, isoindole, cumarine, cumaronecinoline, isochinoline, cinnoline and the like.

Hetero atoms" are atoms other than carbon and hydrogen, preferably nitrogen (N), oxygen (O) or sulfur (S).


The heterocyclic moiety may form a linker L^2 in that the heterocyclic moiety is covalently linked to the phosphor acid ester and the group A.

Likewise, the heterocyclic moiety may form a linker L^1 in that the heterocyclic moiety is covalently linked to the phosphor acid ester and the colloid-active compound X.

In a preferred embodiment of the present invention linker L^1 is selected from the group consisting of a single bond and a C_1 - C_{10} alkandiyl, preferably a C_2 - C_6 -

alkandiyl, especially ethan-1,2-diyl (ethylene) or propan-1,2-diyl or propan-1,3-diyl.


In an especially preferred embodiment linker L^2 is selected from the group consisting of a single bond, a moiety of formula (VII), a moiety of formula (VIII) and a moiety of formula (IX)



wherein the R^3 , R^4 , R^2 , R^5 , R^6 and R^7 are as defined above.

According to an especially preferred embodiment of the invention X is hydroxyethylstarch and said medicinally active substance A is 5-fluorouracil (5-fluoro-1H-pyrimidine-2,4-dione) or a derivative or a prodrug thereof. Alternatively, substance A may be 5-fluorouridine or a 5-fluorouridine derivative.


An especially preferred embodiment of the invention is represented by the formula (V)




wherein HES is hydroxyethylstarch and L^1 , L^2 and n are define as above.

Especially, the phosphoric acid forms an ester linkage with the hydroxethyl group of the hydroxyethyl starch. In this case linker L^1 in formula (V) is a single bond.

According to a further aspect of the invention the compound of the present invention is represented by the following formula (Ia):



wherein n is as defined above and Z is selected from the group of following moieties with formulae Ib to Ig:



wherein

c is an integer ranging from 2 to 12, preferably 3 to 8, more preferably 3 to 6;

d is an integer ranging from 1 to 24, preferably 2 to 18, more preferably 3 to 12;

e is 1 or 2;

R^3 , R^4 , R^2 , R^5 , R^6 and R^7 as well as A are as defined above.


A further embodiment of the invention is a pharmaceutical formulation comprising the compound of the invention.

The pharmaceutical formulation is preferably aqueous and injectable.

It has surprisingly found that the compound of the invention can be obtained by reacting a phosphoramidite, which is linked to a medicinally active substance A or a fluorescence marker A, with a colloid active compound which comprises one or more hydroxyl groups.


The reaction sequence provides an access to pharmaceutically acceptable colloids which are linked with a medicinally active substance or a fluorescence marker and which can be cleaved enzymatically.


Therefore, a further embodiment of the invention is a process for preparing a compound of general formula (I) of the invention by linking a phosphoramidite covalently bonded to a A-L²-group, preferably by linking a compound of formula (VI)








with a compound X-L¹-OH or X which comprise at least one hydroxyl group and subsequently oxidizing and hydrolyzing the linked product so as to form a phosphoric acid diester. The groups X, A, L¹ and L² are as defined above.

In a preferred embodiment of the invention the colloid active compound X, which is preferably a starch, comprises one or more substituents selected from the group consisting of the moieties reflected in the following formulae:







wherein

f is an integer ranging from 1 to 4;

g is an integer ranging from 1 to 10, preferably 1 to 4;

h is an integer ranging from 1 to 4; preferably 2 to 4;

i is an integer ranging from 1 to 4; preferably 2 to 4;

j is an integer ranging from 1 to 4; preferably 2 to 4;

and G is independently selected from -O- or NR¹¹, wherein R¹¹ is selected from H

or C₁ to C₄ alkyl.

The preparation of phosphoramidites is generally known to the person skilled in the art.

A phosphoramidite (RO)₂PNR₂ is a monoamide of a phosphite diester. The key feature of phosphoramidites is their markedly high reactivity towards nucleophiles catalyzed by weak acids e.c., triethylammonium chloride or 1*H*-tetrazole. In these reactions, the incoming nucleophile replaces the NR₂ moiety.

Phosphoramidites derived from protected nucleosides are referred to as nucleoside phosphoramidites and are widely used in chemical synthesis of DNA, RNA, and other nucleic acids and their analogs.

Nucleoside phosphoramidites are derivatives of natural or synthetic nucleosides. They are used to synthesize oligonucleotides, relatively short fragments of nucleic acid and their analogs. Nucleoside phosphoramidites were first introduced in 1981 by Beaucage and Caruthers (Beaucage, S.L.; Caruthers M.H. (1981). "Deoxynucleoside phosphoramidites—A new class of key intermediates for deoxypolynucleotide synthesis". *Tetrahedron Letters* **22**: 1859–1862). In order to avoid undesired side reactions, reactive hydroxy and exocyclic amino groups present in natural or synthetic nucleosides are appropriately protected. As long as a nucleoside analog contains at least one hydroxy group, the use of the appropriate protecting strategy allows one to convert that to the respective phosphoramidite and to incorporate the latter into synthetic nucleic acids. There are three main methods for the preparation of nucleoside phosphoramidites.

The method used most commonly consist in the treatment of a protected nucleoside bearing a single free hydroxy group with phosphorodiamidite under the catalytic action of a weak acid (Nielsen, J.; Marugg, J. E.; Taagaard, M.; Van Boom, J. H.; Dahl, O. (1986). "Polymer-supported synthesis of deoxyoligonucleotides using in situ prepared deoxynucleoside 2-cyanoethyl phosphoramidites".

Rec. Trav. Chim. Pays-Bas **105** (1): 33–34; Nielsen, J.; Taagaard, M.; Marugg, J. E.; Van Boom, J. H.; Dahl, O. (1986). "Application of 2-cyanoethyl N,N,N',N'-tetraisopropylphosphorodiamidite for in situ preparation of deoxyribonucleoside phosphoramidites and their use in polymer-supported synthesis of oligodeoxyribonucleotides". *Nucl. Acids Res.* **14** (18): 7391–7403).

2-cyanoethyl N,N,N',N'-tetraisopropylphosphorodiamidite, the amidite used for the preparation of commercial nucleoside phosphoramidites, is relatively stable. It can be synthesized using a two-step, one-pot procedure and purified by distillation.

According to an alternative method, the protected nucleoside is treated with the phosphorochloridite in the presence of an organic base, most commonly N-ethyl-N,N-diisopropylamine (Hünig's base).

In a further alternative synthetic route for the preparation of phosphoramidites the protected nucleoside is first treated with chloro N,N,N',N'-tetraisopropyl phosphorodiamidite in the presence of an organic base, most commonly N-ethyl-N,N-diisopropylamine (Hünig's base) to form a protected nucleoside diamidite. The latter is treated with an alcohol respective to the desired phosphite protecting group, for instance, 2-cyanoethanol, in the presence of a weak acid.

Nucleoside phosphoramidites can be purified by column chromatography on for example silica gel. To warrant the stability of the phosphoramidite moiety, it is advisable to equilibrate the column with an eluent containing 3 to 5wt.-% of triethylamine and maintain this concentration in the eluent throughout the entire course of the separation.

It has surprisingly found that phosphoramidites can undergo a phosphoramidite coupling reaction with the colloid active compound X which comprises at least one hydroxy group. The reaction can be carried out in the presence of an acidic azole catalyst, 1*H*-tetrazole, 2-ethylthiotetrazole, 2-benzylthiotetrazole, 4,5-dicyanoimidazole, or a number of similar compounds.


Phosphoramidites are readily oxidized with weak oxidating reagents, for instance, with aqueous iodine in the presence of weak bases or with hydrogen peroxide to form the respective phosphoramidates.

According to a preferred embodiment of the present invention the oxidation step is carried out in the presence of a CSO oxidizer, preferably ((1S)-(+)(10-comphorsulfonyl)-oxaziridine).

The naturally occurring nucleotides (nucleoside-3'- or 5'-phosphates) and their phosphodiester analogs are insufficiently reactive to afford an expedite synthetic preparation of oligonucleotides in high yields. The selectivity and the rate of the formation of internucleosidic linkages are dramatically improved by using 3'-O-(*N,N*-diisopropyl phosphoramidite) derivatives of nucleosides (nucleoside phosphoramidites) that serve as building blocks in phosphite triester methodology. To prevent undesired side reactions, all other functional groups present in nucleosides should be rendered unreactive (protected) by attaching protecting groups.

Thus, in a preferred embodiment, groups present at group A which may react with the phosphoramidite such as nucleophilic groups, e.g. OH groups are protected by protecting groups which can be cleaved after the coupling reaction with the colloid active compound occurred.

Scheme 3 shows in an exemplary embodiment a synthetic route to obtain a phosphor acid diester attached to a hydroxyethyl starch.




Scheme 3

Examples

It has been found that medicinally active substances or fluorescence marker can be coupled to colloid active compounds comprising at least one hydroxy group, such as hydroxyethyl starch (HES). In Scheme 3 a reaction sequence is shown for a corresponding 5-fluorouridine derivative. For this purpose, 5-fluorouridine is first lipophilized at the O-2',3' position by a cyclic ketal moiety; the ring size is variable. The ketal moiety can be for example obtained by the synthesis disclosed in E. Malecki, H. Rosemeyer, *Helv. Chim. Acta* **2010**, 93, 1500. An


example is shown in the following formula representing compound **6** wherein $n=14$.



The ketal moiety further acts as a protecting group. A phosphoramidite **2** with $n=14$ (hereinafter **2a**) can be prepared as follows:

Preparation of phosphoramidite **2a**:

5-Fluoro-1-[(4'R,6'R)-2',3',4',5'-tetrahydro-6'-(hydroxymethyl)spiro[cyclopentadecane-1,2'-furo[3,4-d][1,3]dioxol]-4'-yl]pyrimidine-2,4(1H,3H)-dione 2-Cyanoethyldiisopropylphosphoramidite (2a).




Anhydrous compound **6** (256 mg, 0.45 mmol) was 5'-phosphitylated under an nitrogen atmosphere using ethyldiisopropylamine (*Hünig's* base, 147 μ l, 0.85 mmol) and (chloro)(2-cyanoethoxy)(diisopropylamino)phosphine (181 μ l, 0.80

mmol) The reaction mixture was stirred for 15 min at room temperature, and then an ice-cold 5% aq. NaHCO₃ solution (12 ml) was added. The mixture was extracted three times with cold CH₂Cl₂, the combined organic layers were dried (Na₂SO₄), filtered and evaporated on a rotary evaporator (bath temperature, 25°C). Chromatography (silica gel, column: 2 x 8 cm, CH₂Cl₂/MeOH, 8:2, v/v) gave one main zone from which compound **2a** (208 mg, 60 %) was obtained as colourless oil. TLC (CH₂Cl₂/MeOH, 8:2, v/v): *R_f*, 0.95. ³¹P-NMR (CDCl₃): 149.56, 149.41.

From Scheme 3 it can be seen that the synthesis of the end product **3** implies – after the coupling of the P(III) derivative **2** to HES – the oxidation of **2** with I₂/H₂O, followed by a cleavage of the cyanoethyl protecting group in concentrate aqueous ammonia. The reactions are preferably performed under strict exclusion of moisture (Ar atmosphere) and using thoroughly dried hydroxyethyl starch (HES). The advantage of a product such as compound **3** is its enzymatic cleavability by phosphodiesterases.

Scheme 4 shows another route for the preparation of a compound of the invention.

Phosphoramidite **4** which is based on 5-(propyn-1-yl)-2'-deoxyuridine a virostatic compound, active against *Herpes simplex* viruses (A.L. Andronova et al. *Russ. J. Bioorg. Chem.* **2003**, 29, 262-266) has been coupled to HES.




Scheme 4

HES-40 (1 g) was dried by repeated (7 times) lyophilisation (freeze drying) from anhydrous acetonitrile (MeCN) (20 ml, each) during one week. The HES (500 mg) was transferred into a reactor and purged with anhydrous MeCN (10 ml).

Subsequently, the solid was purged with a 0.25 M 4,5-dicyano-imidazole (DCI) activator solution in MeCN (10 ml). Simultaneously, 5-(propin-1-yl)-2'-deoxyuridine 2-(cyanoethyl)(diisopropyl)phosphoramidite (**4**, 1 g) was dissolved in the DCI activator solution and diluted with MeCN to a total volume of 10 ml. After injection into the reactor, the suspension was slightly agitated for 10 min at room temperature. Then, the product was washed twice with anhydrous MeCN (10 ml, each), and, next, an oxidizer solution (0.02 M I₂, THF, pyridine, 10 ml) was injected. After 1 min of agitation within the reactor, the material was washed 4 times with anhydrous MeCN (20 ml, each). For drying N₂ gas was purged through the reactor for several minutes, and the material was removed from the column. ³¹P-NMR ((D₆)DMSO: -1.25 ppm.

CLAIMS

1. A compound of formula (I)



wherein

X is a colloid-active compound;

L¹ is a first linker by means of which X and the phosphate group are covalently linked together;

L² is a second linker by means of which the phosphate group and A are covalently linked together;

A is a medically active substance or a fluorescence marker;

n is an integer of at least 1.

2. The compound according to claim 1, characterized in that said colloid-active compound X is selected from the group consisting of amyloses, amylopectins, acemannans, arabinogalactans, galactomannans, galactoglucomannans, xanthans, carrageenan, hyaluronic acid, deacetylated hyaluronic acid, starch and modified starch.

3. The compound according to claim 2, characterized in that said modified starch is selected from the group consisting of hydroxyalkyl starches, esterified starches, carboxyalkyl starches, hydroxyalkyl carboxyalkyl starch, aminated hydroxyalkyl starch, aminated hydroxyalkyl carboxyalkyl starch and aminated carboxyalkyl starch.


4. The compound according to claim 2, characterized in that said modified starch is selected from hydroxyethyl starch or aminated hydroxyethyl starch.



5. The compound according to claims 1 to 4, characterized in that said colloid-active compound has an average molecular weight of from 20,000 to 800,000 daltons, preferably from 25,000 to 500,000 daltons, especially from 30,000 to 200,000 daltons.

6. The compound according to at least one of claims 1 to 5, characterized in that the degree of substitution, DS, of the modified starch, especially hydroxyethyl starch, is from 0.2 to 0.8, preferably from 0.3 to 0.6.

7. The compound according to at least one of claims 1 to 6, characterized in that said medicinally active compound A is selected from the group consisting of antibiotics, chemotherapeutics, cytostatic agents, antigens, oligonucleotides, mediators, false metabolic substrates, analgetics and cytotoxic substances.

8. The compound according to at least one of claims 1 to 7, characterized in that said medicinally active compound A is selected from formula (II) to (IV)





wherein

R^1 is H or an organic moiety, preferably a C_1 - C_{28} chain which may be branched or linear and which may be saturated or unsaturated and which may optionally be interrupted and/or substituted by one or more hetero atom(s) (Het1) and/or functional group(s)(G1); or

R¹ is a C₃-C₂₈ moiety which comprises at least one cyclic structure and which may be saturated or unsaturated and which may optionally be interrupted and/or substituted by one or more hetero atom(s) (Het1) and functional group(s)(G1);

R² is H or an organic moiety comprising 1 to 30 carbon atoms;

R³ and R⁴ represent independently from each other H or a C₁-C₂₈-alkyl moiety which may optionally be substituted or interrupted by one or more heteroatom(s) and/or functional group(s); or

R³ and R⁴ form a ring having at least 5 members, preferably a ring having 5 to 8 carbon atoms and wherein the ring may be substituted or interrupted by one or more hetero atom(s) and/or functional group(s);

R⁵ and R⁶ represent independently from each other H or a C₁-C₂₈-alkyl moiety which may optionally be substituted or interrupted by one or more heteroatom(s) and/or functional group(s);

or


R⁵ and R⁶ form a ring having at least 5 members, preferably a ring having 5 to 18 carbon atoms and wherein the ring may be substituted or interrupted by one or more hetero atom(s) and/or functional group(s);

R⁷ is a hydrogen atom or -O-R⁸;

R⁸ is H or C₁-C₂₈ chain which may be branched or linear and which may be saturated or unsaturated and which may optionally be interrupted and/or substituted by one or more hetero atom(s) (Het1) and/or functional group(s)(G1).

9. Compound according to claim 8 wherein

R¹ is selected from H,



substituted or unsubstituted cyclic terpene moieties,

wherein

R^9 and R^{10} are independently selected from C_1 to C_{30} alkyl,


b is an integer ranging 1 to 4, preferably n is 1 or 2; and

a is an integer ranging from 1 to 20, preferably 2 to 18

10. The compound according to at least one of claims 1 to 7, characterized in that said fluorescence marker is selected from the group consisting of fluorescein isothiocyanate (FITC), phycoerythrin, rhodamide, 2-aminopyridine and coumarine dyes.


11. The compound according to claim 1, wherein said X is hydroxyethylstarch and said medicinally active substance A is 5-fluorouracil (5-fluoro-1*H*-pyrimidine-2,4-dione) or a derivative or a prodrug thereof.

12. The compound according to claim 1 which is represented by the formula (V)



wherein HES is hydroxyethylstarch.

13. A pharmaceutical formulation comprising the compound according to at least one of claims 1 to 12.
14. The pharmaceutical formulation according to claim 13, characterized in that said formulation is aqueous and injectable.
15. A process for preparing a compound of general formula (I) according to at least one of claims 1 to 12 by linking a compound of formula (VI)



with a compound X-L¹-OH or X which comprise at least one hydroxyl group and subsequently oxidizing and hydrolyzing the linked product so as to form a phosphor acid diester.

Colloid bonded Medicinal Compounds

Abstract

The invention relates to colloids bound medicinal compounds or fluorescent markers, to a process for the preparation thereof, and to a pharmaceutical formulation containing such compounds.